

3050 Spruce Street Saint Louis, Missouri 63103 USA Telephone 800-325-5832 • (314) 771-5765 Fax (314) 286-7828 email: techserv@sial.com sigma-aldrich.com

# **ProductInformation**

## **Xanthine** Sigma Ultra

Product Number X 4002 Store at Room Temperature

#### **Product Description**

Molecular Formula: C<sub>5</sub>H<sub>4</sub>N<sub>4</sub>O<sub>2</sub> Molecular Weight: 152.1 CAS Number: 69-89-6

pK<sub>a</sub>: 7.7, 11.9<sup>1</sup>

 $\lambda_{\text{max}} = 277 \text{ nm } (\text{pH} = 10)^1$ Extinction Coefficient:  $E^{\text{mM}} = 9.3 \text{ (pH} = 10)^1$ 

Xanthine is a catabolic product of purine nucleotide metabolism. It is degraded by xanthine oxidase to uric acid. Natural sources of xanthine or its derivatives include animal organs, yeast, potatoes, coffee beans, and tea.<sup>2</sup> Xanthine and xanthine oxidase are used to generate superoxide radicals used to measure the activity of superoxide dismutase.3

Trace elemental analyses have been performed on the SigmaUltra xanthine. The certificate of analysis provides lot specific results. SigmaUltra xanthine is for applications, which require tight control of elemental content.

#### **Precautions and Disclaimer**

For Laboratory Use Only. Not for drug, household or other uses.

### **Preparation Instructions**

Xanthine is soluble in sodium hydroxide solutions and in acidic solutions. It is soluble in 1 M NaOH (50 mg/ml), with sonication for less than 5 minutes, yielding a clear solution. It is slightly soluble in water (1 g/14.5 L, 16 °C) and in ethanol.2

## Storage/Stability

The decomposition of solutions in 0.5 M H<sub>2</sub>SO<sub>4</sub> or in 10 M NaOH is less than 10% after one hour at 100 °C.4

Stock solutions of at least 10 mM in sodium hydroxide can be stored at 2-8 °C for one week.5

#### References

- 1. Specifications and Criteria for Biochemical Compounds, 3rd ed., National Academy of Sciences (Washington, DC: 1972), p.182.
- The Merck Index, 13th ed., Entry# 10116.
- McCord, J. M., and Fridovich, I., Superoxide Dismutase. An Enzymic Function For Erythrocuprein (Hemocuprein). J. Biol. Chem., **244(22)**, 6049-6055 (1969).
- 4. Data for Biochemical Research, 3rd ed., Dawson, R. M. C., et al., Oxford University Press (New York, NY: 1986), p. 94-95.
- 5. Heinz, F., and Reckel, S., in Methods Of Enzymatic Analysis, 3rd ed., Vol. 3, Bergmeyer, H. U., ed., Academic Press (New York, NY: 1983), p. 210-216.

ARO/RXR 11/02